

Kinetics and thermodynamics in the protein-ligand interactions during activity in the bifunctional FAD synthetase from *Corynebacterium ammoniagenes*

P01-01

M. Sebastián^I, A. Serrano^I, A. Velázquez-Campoy^I, M. MEDINA^I

^I*Departamento de Bioquímica y Biología Molecular y Celular, Facultad de Ciencias, and Institute of Biocomputation and Physics of Complex Systems, Universidad de Zaragoza, Zaragoza, Spain*

Prokaryotic bifunctional FAD synthetases (FADSs) are bimodular enzymes exhibiting ATP:riboflavin kinase (RFK) activity in its C-terminal module and FMN:ATP adenylyltransferase (FMNAT) activity in its N-terminal. These activities provide the organism with the flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) cofactors. The RFK activity of FADS from *Corynebacterium ammoniagenes* (CaFADS) has a strong inhibition at mild concentrations of the RF substrate. Selective inhibition of key enzymes is a common tool to regulate metabolic pathways. Since FMN and FAD act as cofactors in a plethora of flavoproteins and flavoenzymes in all living organisms, inhibition of the RFK activity in some family members might contribute to the flavin cellular homeostasis, and, therefore, it is a topic worthy of study. Here we use a truncated CaFADS variant that only contains the C-terminal RFK module, being it similarly functional in the RFK activity as the full length enzyme. The steady-state characterization of this variant indicates that besides inhibition by the RF substrate, both of the reaction products, ADP and FMN, also inhibit the RFK activity. The use of pre-steady-state kinetics collectively with isothermal titration calorimetry allows us to present a kinetic and thermodynamic explanation of such inhibitory behavior related to ligand binding that is coherent with the structural conformational changes occurring during the RFK catalysis in CaFADS. Furthermore, these methods can be also used to evaluate the behavior of the full-length enzyme.